



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\begin{aligned}
 &= \frac{8}{\pi^3} \int_0^{\frac{\pi}{2}} \int_{\frac{1}{2}\pi-\theta}^{\frac{1}{2}\pi} \varphi(\pi-\varphi) \operatorname{cosec}^2 \theta d\theta \sin \varphi \cos \varphi d\varphi \\
 &= \frac{1}{\pi^3} \int_0^{\frac{\pi}{2}} (\pi^2 - 4\theta^2 + 2\theta^2 \operatorname{cosec}^2 \theta - 4\theta \cot \theta + 2) d\theta \\
 &= \frac{1}{3} + \frac{1}{\pi^2}.
 \end{aligned}$$

QUERY. "In Thompson and Tait's 'Elements of Natural Philosophy' it is stated that, 'at the southern base of a hemispherical hill radius a and density ρ , the true latitude (as measured by the aid of the plumb-line) is diminished by the attraction of the mountain by the angle $\frac{2}{3}\rho\pi a \div (G - \frac{4}{3}\rho a)$, where G = the attraction of the earth in same units.' How is this proved?"

ANSWER BY HENRY HEATON.

As the angle is so small that it is practically equal to its tangent, if G = attraction of the earth, $\frac{2}{3}\rho\pi a$ = horizontal, and $\frac{4}{3}\rho a$ = vertical attraction of the hill, then it is evident that the angle of deflection of the plumb-line = $\frac{2}{3}\rho\pi a \div (G - \frac{4}{3}\rho a)$.

Where the unit of attraction is the attraction of a mass whose volume is 1, density 1, and distance 1, it has been shown, page 194, Vol. II, of the ANALYST, that the horizontal attraction of the hill = $\frac{2}{3}\rho\pi a$; and that the vertical attraction = $\frac{4}{3}\rho a$, I shall now proceed to show.

Taking the origin of co-ordinates at the center of the spherical surface, the attraction exerted by an element of the mass resolved in a vertical direction = $\frac{\rho dx dy dz}{[(x+a)^2 + y^2 + z^2]^{\frac{3}{2}}}$. Hence the vertical attraction of the hill is

$$\begin{aligned}
 &\rho \int_{-a}^{+a} \int_{-u}^{+u} \int_0^v \frac{dx dy dz}{[(x+a)^2 + y^2 + z^2]^{\frac{3}{2}}} = \rho \int_{-a}^{+a} \int_{-u}^{+u} \left[\frac{dx dy}{[(x+a)^2 + y^2]^{\frac{1}{2}}} \right. \\
 &\quad \left. - \frac{dx dy}{[2a^2 + 2a]^{\frac{1}{2}}} \right] = \int_{-a}^{+a} \left[\log \left(\frac{\sqrt{(2a)} + \sqrt{(a-x)}}{\sqrt{(2a)} - \sqrt{(a-x)}} \right) dx - \frac{2\sqrt{(a-x)} dx}{\sqrt{(2a)}} \right] \\
 &= \frac{4}{3}\rho a; \text{ where the limits of integration, } u \text{ and } v, \text{ are, respectively, } u = \sqrt{(a^2 - x^2)} \text{ and } v = \sqrt{(a^2 - x^2 - y^2)}.
 \end{aligned}$$

If we put r for radius of the earth, and δ , its mean density, $G = \frac{4}{3}\delta r$. Substituting this for G in the expression for the angle of deflection, we get, $\rho\pi a \div 2(\delta\pi r - \rho a)$. As a is very small in comparison with r the expression for the deflection is, practically, $\rho a \div 2\delta r$.

[Mr. Adcock obtains precisely the same result as the last above written, viz.; $\rho a \div 2\delta r$.]